

Poster presentation

Influence of external input on waxing and waning of neuronal network oscillations

Oscar J Avella Gonzalez^{*1}, Ronald van Elburg², Huibert Mansvelder¹, Jaap van Pelt¹ and Arjen van Ooyen¹

Address: ¹Department of Integrative Neurophysiology, VU University Amsterdam, 1081 HV The Netherlands and ²Department of Artificial Intelligence, University of Groningen, 9700 AB, The Netherlands

Email: Oscar J Avella Gonzalez^{*} - oscar.avella@cncr.vu.nl

^{*} Corresponding author

from Eighteenth Annual Computational Neuroscience Meeting: CNS*2009
Berlin, Germany. 18–23 July 2009

Published: 13 July 2009

BMC Neuroscience 2009, **10**(Suppl 1):P257 doi:10.1186/1471-2202-10-S1-P257

This abstract is available from: <http://www.biomedcentral.com/1471-2202/10/S1/P257>

© 2009 Gonzalez et al; licensee BioMed Central Ltd.

Introduction

Experimental observations have reported modulation of cortical oscillations as phases of high synchronization (waxing) followed by periods of reduced synchronization (waning) [1-3]. Although the phenomenon is present in almost all frequency bands, it is still not understood how this is driven. Here we study whether this phenomenon can occur in a network of inhibitory (I) and excitatory (E) cells and what effect external inputs have.

Methods

Using NEURON, we model a network of N_e excitatory and N_i inhibitory cells such that $N_e/N_i = 4$. The cells have a single compartment, and include passive channels and voltage dependent Na^+ , K^+ channels. Synaptic connections are random, projecting GABA synapses from I to I and I to E cells and AMPA synapses from E to E and E to I cells. To stimulate the network, each cell receives a baseline of current and a stream of spikes delivered at random intervals across the simulated period.




Figure 1
Raster plots of E (top) and I (middle) populations, during waxing and waning of a beta oscillation, and firing rate histograms (bottom).




Figure 2
Wavelet transform of the activity in the E population for the same time period as shown in figure 1.

Results

We show that in a stable oscillatory network, waxing and waning occurs without the need for other synaptic mechanisms than the spike generating K^+ and Na^+ channels. The phenomenon can be modulated by changing the characteristics of the external input, such as number of spikes, mean inter-spike interval, randomness and whether E or I cells receive the external input. See figures 1 and 2.

Acknowledgements

The work was supported by the EU grant MC-RTN NEURoVERS-it (019247).

References

1. Hughes SW, Loerincz M, Cope DW, Blethyn KL, Kékesi KA, Parri HR, Juhász G, Crunelli V: **Synchronized oscillations at α and β frequencies in the lateral geniculate nucleus.** *Neuron* 2004, **42**:253-268.
2. Palva S, Palva JM: **New vistas for α -frequency band oscillations.** *Trends Neurosci* 2007, **30**:150-158.
3. Courtemanche R, Lamarre Y: **Local field potential oscillations in primate cerebellar cortex: synchronization with cerebral cortex during active and passive expectancy.** *J Neurophysiol* 2005, **93**:2039-2052.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

